

These standards are to be applied on individual truck loads and must not be averaged over a number of loads.
Segregations indicated on this chart are only available where announced.

		Compass Minatour Zena CL Cyclops Neo CL La Trobe Commodus CL Maximus CL RGT Planet Spartacus CL All Other Varieties	MALTING BARLEY			BARLEY	
			CM1 MN1 ZE1 CY1 NE1 LA1 CD1 MA1 PL1 SP1 -	CM2 MN2 ZE2 CY2 NE2 LA2 CD2 MA2 PL2 SP2 -	CM3 MN3 ZE3 CY3 NE3 LA3 CD3 MA3 PL3 SP3 -	BAR1 BAR1 BAR1 BAR1 BAR1 BAR1 BAR1 BAR1 BAR1 BAR1	BAR2 BAR2 BAR2 BAR2 BAR2 BAR2 BAR2 BAR2 BAR2 BAR2
BINNED GRADE							
All malting varieties must be segregated and of the current season							
Test Code							
VARP	Varietal Purity minimum (% by count per 100 Grain Sample or approved method) All approved 2 row malting varieties of the current season	95	95	95	All 2 row varieties		
MOGR	Moisture maximum (%)	12.5	12.5	12.5	12.5	12.5	
PRGR	Protein minimum (%)	9.0	9.0	9.0	No Limit	No Limit	
	Protein maximum (%)	12.0	12.0	12.8	No Limit	No Limit	
TWT	Test Weight minimum (kg/hl)	65.0	65.0	65.0	62.5	60.0	
RET	Retention minimum (% by weight) Above 2.5mm Agtator screen	70.0	62.0	58.0	No Limit	No Limit	
SCRN	Screenings maximum (% by weight) Below 2.2mm Agtator screen	7.0	10.0	No Limit	15.0	25.0	
GE72	Germination	95	95	95	No Limit	No Limit	
FALL	Falling Number minimum (seconds)	300	300	300	N/A	N/A	
Defective Grains – Maximum tolerance as per method specified. Note: NIL tolerance applies to the entire load.							
Defective definitions are to be read in conjunction with the photo in the Visual Recognition Standards Guide which depicts the minimum standard for a grain to be classified as defective.							
SHOT	Shot maximum (% by count per 100 Grain Sample) - Note: The NIL tolerance for shot grains does not apply if a Falling Number analysis is conducted.	NIL	NIL	NIL	No Limit	No Limit	
SPRO	Sprouted maximum (% by count per 100 Grain Sample) - Any visible evidence of the shoot or root system beginning to emerge from the germ. Note: NIL tolerance for sprouted grains does not apply to malt grades if a Falling Number analysis is conducted.	NIL	NIL	NIL	NIL	5	
BTIP	Dark Tipped (% by count per 100 Grain Sample)	10	10	10	No Limit	No Limit	
FFUN	Field Fungi maximum (% by count per 100 Grain Sample) Seed coat has the appearance of black spotting occurring anywhere on the grain. Coverage greater than approximately 10% of the grain surface is considered defective, otherwise classified as sound. Grey surface discolouration of the kernel. Does not include Severely Damaged.	5	5	5	No Limit	No Limit	
SKIN	Skinnings maximum (% by count per 100 Grain Sample)	15	15	15	No Limit	No Limit	
CLD	Cleaved maximum (% by count per 100 Grain Sample) - A split on any part of the grain that exposes the white endosperm.	1	1	1	No Limit	No Limit	
DIST	Distorted maximum (% by count per 100 Grain Sample) - Grain which is collapsed on the dorsal side. Grains may appear orange in colour.	5	5	5	10	10	
GREE	Dry Green or Sappy maximum (% by count per 100 Grain Sample)	1	1	1	No Limit	No Limit	
PFUN	Pink Fungal Maximum (count per half litre)	1	1	1	20	30	
DAMI	Insect Damaged maximum (count per half litre)	10	10	10	85	85	
SEVE	Severely Damaged maximum (count per half litre above 2.2mm and 2.5mm screens) Mould, heat damaged / burnt, diseased or other serious visual defects. Grains that have become severely discoloured. Grains appear dark brown, or in severe cases, blackened. May also appear discoloured under the husk on the kernel. Does not include Field Fungi.	1	1	1	2	5	
BKN	Broken maximum (% by weight per 100 gram Sample)	2	2	2	5	5	
Foreign Seed Contaminants – Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per half litre.							
Except TYPE(1) in which the maximum applies on an individual seed basis per half litre.							
WS1	TYPE(1): Colocynthis, Poppy (Field, Horned), Jute, Long Head Poppy, Mexican Poppy, New Zealand Spinach, Parthenium Weed** (QLD Only), Saffron Thistle, Wild Poppy	8*	8*	8*	8*	8*	
WS2	TYPE(2): Castor Oil Plant, Coriander, Crow Garlic/ Wild Garlic, Darling Pea, Opium Poppy, Parthenium Weed** (NSW/VIC/SA), Peanut seeds and pods, Ragweed, Rattlepods, Starburr, St. John's Wort	NIL	NIL	NIL	NIL	NIL	
WS3A	TYPE(3a): Bathurst Burr, Bellvine, Branched Broomrape, Bulls Head/Caltrop/Cats Head, Cape Tulip, Cottonseed, Dodder, Noogoora Burr, Thornapple	2	2	2	2	2	
WS3B	TYPE(3b): Vetch (Commercial), Vetch (Tare)	4	4	4	4	10	
WS3C	TYPE(3c): Heliotrope (Blue), Heliotrope (Common) *****Note: included in this Type are tolerances for seeds or pods.	1 Pod / 4 Seeds	1 Pod / 4 Seeds	1 Pod / 4 Seeds	1 Pod / 4 Seeds	1 Pod / 4 Seeds	
WS3D	TYPE(3d): Double Gees / Spiny Emex / Three Cornered Jack	1	1	1	1	1	
WS4	TYPE(4): Bindweed (Field), Cutleaf Mignonette seeds, Darnel (Drake Seed), Hexham Scent/King Island Melilot*** (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshade, Paddy Melon, Skeleton Weed, Variegated Thistle	20	20	20	20	20	
WS5	TYPE(5): Knapweed (Creeping/Russian), Sesbania Pea, Patterson's Curse/ Salvation Jane	40	40	40	40	40	
WS6	TYPE(6): Colombus Grass, Johnson Grass ****	NIL	NIL	NIL	40	40	
WS7A	TYPE(7a): Adzuki Beans, Broad Beans, Chickpeas, Clover, Corn (Maize), Cowpea, Faba Beans, Lentils, Lupins, Peas (Field), Medic Pods, Safflower, Soybean, Sunflower and any other seeds or pods greater than 5mm in diameter including broad bean. Includes Onion Weed Pods regardless of size	1	1	1	10	20	
WS7B	TYPE(7b): 6 row barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Musk Weed, Onion Weed, Phalaris Glumes, Poverty Weed, Ryegrass on stalk, Sheep Weed, Sorghum Grain, Three Horn Bedstraw, Turnip Weed Pod (regardless of size) and any other foreign seed not specified in Types 1-7(a), Foreign Grain (GC9A), Variation A (GC9B), Variation B (GC9C) or in SFS.	50	50	50	150	300	
GC9A	Foreign Grain: Wheat, Cereal Rye, Triticale, Rice, Cultivated Oats (Common)	85	85	85	500	1500	
GC9B	Variation A: Oats (Black or Wild)***** , Wild Radish Pods	25	25	25	50	100	
GC9C	Variation B: Barley with Coloured Aleurone Layer (Blue or Black). Also includes black hulled varieties.	NIL	NIL	NIL	100	100	
SFS	Small Foreign Seeds maximum (% by weight) Foreign seeds not already specified in Types 1-9(c) that fall below the 2.2mm screen during the screenings process.	0.6%	0.6%	0.6%	1.2%	2.0%	
* Individual Seed Basis **Parthenium Weed is a NIL tolerance in NSW/VIC/SA *** Hexham Scent is only acceptable if no tainting odour is present **** TYPE(6) limit in QLD is 40 per half litre for all barley grades ***** Heliotrope pods must be opened and the seeds counted *****Counted individually if clustered							
Other Contaminants – Maximum tolerance as per method specified. Note: NIL tolerance applies to the entire load except where stated otherwise.							
All Foreign Seed Pods not listed above such as those that are 5mm or less in diameter are included as Foreign Material, whether whole pods or part thereof.							
FORM	Foreign Material maximum (% by weight per half litre) - Other than already specified. Includes: Milk Thistle Pods, pieces of seed pods.	1%	1%	1%	1%	1%	
PICK	Pickling Compounds maximum (entire load)	NIL	NIL	NIL	NIL	NIL	
CHEM	Chemicals Not Approved for Barley maximum (entire load)	NIL	NIL	NIL	NIL	NIL	
ERGC	Cereal Smut/Cereal Ergot maximum (entire load) - Includes: Ball and gall smut, other smut species, cereal ergots.	NIL	NIL	NIL	NIL	NIL	
ERGR	Ryegrass Ergot maximum (length in cm when pieces are aligned per half litre)	0.5cm	0.5cm	0.5cm	0.5cm	0.5cm	
LIVE	Stored Grain Insects and Pea Weevils: Live maximum (entire load)	NIL	NIL	NIL	NIL	NIL	
INLG	Field Insects – Large, dead or alive (count per half litre) - Includes: Desianthra Weevil, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, Ladybirds, Grasshoppers, Locusts and Wood Bugs, whole or parts thereof. These are insect contaminants of grain that do not cause damage to stored grains.	3	3	3	3	3	
INSM	Field Insects – Small, dead or alive (count per half litre) - Includes: all species of Aphid, Minute Mould Beetle, Mites & stored grain insects (dead only). These are insect contaminants of grain that do not cause damage to stored grains.	10	10	10	10	10	
SNAL	Snails: Live or Dead maximum (count per half litre)	1	1	1	1	4	
GUMN	Gumnuts maximum (count per 2.5L) - Whole or pieces of any size and maturity level	1	1	1	1	1	
LSMT	Loose Smut maximum (weight in grams per half litre)	0.1	0.1	0.1	0.1	0.1	
SAND	Sand maximum (count per half litre)	50	50	50	50	50	
EART	Earth maximum (count per half litre) - Defined as a clod of dirt, being 5mm or less in diameter.	3	3	3	3	3	
STON	Stones maximum (grams per 2.5L) - Maximum total weight of all stones retained above the 2.2mm and 2.5mm screens per 2.5L	4.0	4.0	4.0	4.0	4.0	
OBJM	Objectional Material maximum (entire load) Presence of meat meal, blood meal, fish meal, poultry offal meal or other animal proteins. Sticks/Stubble (>3cm in length and 1cm in diameter), glass, concrete, metal, fertiliser, animal excreta, animal carcasses, tainting agents or any other commercially unacceptable contaminant, odour or taste	NIL	NIL	NIL	NIL	NIL	