

THIS CHART IS INTENDED TO ACT AS A GUIDE ONLY

These standards are to be applied on individual truck loads and must not be averaged over a number of loads.

Issued on 01 August 2025

Segregations indicated on this chart are only available where announced. Test Code Binned Grade NIP1 NIP₂ Physical Characteristics - The Lentils shall be red, hard and well filled. Purity minimum (% by weight) Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats of the following varieties - Nipper (L015), PBA Herald XT 97 97 VARP (L019), PBA Hurricane XT (L023), PBA Highland XT (L029), GIA Lightning (L032), GIA Thunder (L034) MOGR | Moisture maximum (%) 14 14 Defective Grains - maximum % by weight per minimum 200g sample. Defective definitions are to be read in conjunction with the photo in the Visual Recognition Standards Guide which depicts the minimum standard for a grain to be classified as defective. Total Defective maximum (% by weight per minimum 200g sample) Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, and all Red Lentil seed material falling through the slotted 2.0mm DEFG 4 8 OF WHICH (Included in Total Defective) 1 1 Variety (% by weight per minimum 200g sample) Red Lentils not of the specified varieties. **OF WHICH (Included in Total Defective)** Poor Seed Coat Colour maximum (% by weight per minimum 200g sample) Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any 1 3 disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta) Does not include Contrasting Colour. Where any poor colour is present on the seed coat, it is recommended the kernel also be inspected OF WHICH (Included in Total Defective)
Poor Kernel Colour (% by weight per minimum 200g sample) Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any 1 disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Severely Damaged maximum (count per 200g sample) 1 1 Mould, heat damaged/burnt, or other serious visual defects. Grains that have become severely discoloured. Not included in Total Defective Contrasting Colour (% by weight per minimum 200g sample)
Contrasting colour of the seed coat in the same variety. Not classified as defective or included in Total Defective. COCO 1 1 Foreign Seed Contaminants - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200g. Except Type(1) in which the maximum applies on an individual seed basis per 200g. TYPE(1): Colocynth, Doublegees / Spiny Emex / Three Cornered Jack, Jute, Long Head Poppy, Mexican Poppy, New Zealand Spinach, Parthenium Weed** (QLD only), Poppy (Field), Poppy (Horned), Wild Poppy 4* 4* TYPE(2): Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Parthenium Weed** (NSW/VIC/SA), Ragweed, NIL NIL WS2 Rattlepods, Starburr, St. Johns Wort WS3A TYPE(3a): Bathurst Burr, Bulls Head/Caltrop/Cats Head, Cape Tulip, Cottonseed, Dodder, Noogoora Burr, Thornapple 1 1 TYPE(3b): Vetch (Tare), Vetch (Commercial) 2 2 WS3C TYPE(3c): Heliotrope (Blue)***, Heliotrope (Common)*** 4 4 TYPE(4): Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent / King Island Meliltot ****, Hoary Cress, Mintweed, 10 10 **WS4** Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle TYPE(5): Knapweed, Sesbania Pea, Pattersons Curse / Salvation Jane 20 20 WS5 TYPE(6): Colombus Grass, Johnson Grass, Saffron Thistle, Clover (Pods), Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed 5 5 (Pods), Sow Thistle (Heads), Wild Radish (Pods), Trefoil (Pods) TYPE(7a): Adzuki Beans, Broad Beans, Cowpea, Chickpeas (Desi / Kabuli), Faba Beans, Lupin, Maize / Corn, Mung beans, Peas (Field), WS7A 10 10 Pigeon Peas, Soybean, any other seeds or pods greater than 5mm diameter not listed in the Standards TYPE(7b) (seeds in total per 200 gram): Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild)****** Oats (Sand)*******, Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, 10 10 WS7B Cereal Seeds(C) or SFS. TYPE(7c): Safflower, Sunflower 1 1 WS7C TYPE(8): Bellvine 100 100 **WSC** Cereals(C): Barley, Corn (Maize), Durum, Oats (Common)******, Rice, Rye (Cereal), Sorghum, Triticale, Wheat 2 2 Small Foreign Seeds maximum (% by weight) **SFS** 0.6% 0.6% Foreign seeds not already specified in Types 1-C that fall in the catchpan during the screenings process *Individual seed basis **Parthenium Weed is a **NIL** tolerance in NSW/VIC/SA seeds counted *****Hexham Scent is only acceptable if no tainting odour is present ******Counted individually if clustered ***Heliotrope pods must be opened and the seeds counted Other Contaminants - Maximum per 200g sample unless otherwise stated. Note: NIL tolerance applies to the entire load. Foreign Material maximum (% by weight) 3 3 **FMAT** Includes: unmillable material and all vegetable matter other than red Lentil seed material, including any Foreign Seeds. OF WHICH UNML 0.5 0.5 Unmillable Material maximum (% by weight) - Includes: soil, stones and non-vegetable matter. OF WHICH 0.3 SOIL 0.3Soil maximum (% by weight) Snails maximum (by count per 200g) 1 1 SNAL Dead or alive. Whole or substantially whole (more than half) including bodies per 200g sample Gumnuts maximum (count per 2 kg sample) 1 1 **GUMN** Whole or pieces of gumnut of any size and maturity level. Field Insects Live or Dead maximum (by count per 200g) 15 15 **GWBL** Includes: ladybirds, minute mould beetles, pea weevil (dead only), sitona weevil, desiantha weevil, wood bugs, other field insects. **GRLO** Grasshoppers & Locusts, Live or Dead maximum (by count per 200g sample) 2 2 Objectionable Material maximum (entire load) Includes: animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, pea weevil **OBJM** NIL NIL (live only), glass, metal, concrete, pickling compounds, tainting agents, odours, fertiliser, sticks or any other commercially unacceptable contaminant, smell or taste. **ERGR** 2cm Ryegrass Ergot maximum (length in cm when pieces are laid end to end per 200g sample) 2cm Non Approved Treatment Chemicals or Treatment Levels above Legal Tolerances maximum (entire load) NIL NIL CHEM Residues of any chemical compound not approved for use on pulses, used in contravention of the labelled instructions or chemicals in excess of the Australian MRL

GrainCorp Enquiries Toll Free 1800 809 482 www.graincorp.com